

# Influence of **dynamics** and **metallicity** on the formation and evolution of **black-hole binaries** in **star clusters**

PhD student: **Brunetto M. Ziosi**

Supervisors: **Dr. Michela Mapelli, Prof. Giuseppe Tormen**

Dept. of Physics and Astronomy G. Galilei (Univ. of Padova),  
INAF-Padova,  
FIRB 2012

PhD Workshop  
Padova, 2013-11-28

# Outline

## 1 Overview

- DCOBs and GWs
- Stellar evolution and BH mass
- Star cluster dynamics

## 2 Tools

- Direct N-body simulations


## 3 Results

- DCOB population
- Orbital properties distribution
- Mass distribution
- Coalescence times

## 4 Conclusions


# Why DCOBs?

- CO: neutron stars and stellar black-holes
- DCO binaries during inspiral and merger events produce **GWs** we could observe in the near future
- Simulations provide theoretical **models** to interpret Advanced **Virgo/LIGO** upcoming data
- Key quantities:
  - **Number** of DCOBs
  - BH **mass** spectrum
  - Binary **orbital properties**



# Why stellar evolution and metallicity (Z)?

- Massive stars **lose mass** by stellar winds
- Winds efficiency **depends** on **metallicity**
- Stars with  $M_{\text{fin}} \geq 40M_{\odot}$  are expected to collapse to a BH **without SN explosion**  
(Fryer 1999, Fryer&Kalogera 2001)
- BHs formed from direct collapse are **more massive** than BHs formed from SN
- Metal-poor stars lose less mass by stellar winds  $\Rightarrow$  more likely to collapse to BH directly




# Why dynamics? Why YSCs?

- **YSCs** are **birthplace** for  $\sim 80\%$  of stars in the local universe (Lada&Lada, 2003)
- (Collisional) YSCs are
  - young ( $< 100$  Myr)
  - relatively massive ( $10^3 - 10^7 M_{\odot}$ ),
  - dense ( $10^3 - 10^6 \star / pc^3$ )

groups of stars


- YSCs are sites of **intense dynamical activity**: central  $t_{\text{relax}} \sim 10 - 100$  Myr




# Why dynamics? Why YSCs?

- **3-body encounters:** close encounters between a single object and a binary
- If kinetic **energy is transferred** from the binary to the single object  $\Rightarrow$  semi-major axis decreases (**hardening**)
- **"Hard" binary:**  $\frac{Gm_1 m_2}{2a} \geq \frac{1}{2} \langle m \rangle \sigma^2$
- Hard binaries tend to become harder as effect of three-body encounters (Heggie 1975)
- The single star can take the place of one of the stars in the binary  $\Rightarrow$  **exchange**
- High probability if  $m_{\text{single}} \gtrsim m_i$  and possibility of high final  $e$ )
- BHs have high masses  $\Rightarrow$  high probability to acquire a companion through 3-body exchange

## Hardening



## Exchange



# Tools


- 200 N-body realizations of the same cluster for each  $Z = 0.01, 0.1, 1Z_{\odot}$
- **StarLab**: Kira (GPU) + SeBa (CPU)  
(Portegies Zwart et al. 2001)
- Our simulations combine **dynamics + up-to-date recipes for Z-dependent stellar evolution**
- Stellar-evolution recipes:
  - accurate **metallicity-dependent** stellar evolution (Hurley et al. 2000) and stellar winds (Vink et al. 2001; Vink & de Koter 2005; Belczynski et al. 2010)
  - the possibility of massive BHs formation by **direct collapse** (Fryer 1999; Fryer & Kalogera 2001; Mapelli et al. 2009)

## 600 simulations

Parameter	Value
$W_0$	5
$N_*$	5500
$M_{\text{tot}}$	$\sim 3500M_{\odot}$
$r_c$ [pc]	0.4
$c = r_t/r_c$	1.03
IMF	Kroupa (2001)
$m_{\text{min}}$ [ $M_{\odot}$ ]	0.1
$m_{\text{max}}$ [ $M_{\odot}$ ]	150
$f_{\text{PB}}$	0.1
$Z$ [ $Z_{\odot}$ ]	0.01, 0.1, 1
Sim. time	100 Myr
MW typical, e.g. Orion Nebula Cluster	

# DCOB population


- DBH distribution
- Mean number of DBHs:  $\sim 3$
- Max number of DBHs: 18
- # NS  $\sim 4$  # BH but
- # DBH  $\sim 10$  # DNS due to **dynamics**
- Negligible **dependence on Z**, but... (see after)



# DBH population in time

Low-Z case vs higher metallicities:

- Build up the DBH population **before** high-Z case
- Higher BH masses allowed  $\Rightarrow$  **earlier** DBHBs **formation**
- Higher BH masses allowed  $\Rightarrow$  more stable binaries & **longer lifetime**




Ziosi et al., in prep

# Orbital properties


- Distributions of orbital parameters at minimum semi-major axis
- Critical for **coalescence** times and mergers **detection**
- SMA and period span a **wide range**
- **DNS** (grey) are 10 times less numerous but have small SMA and short periods

Ziosi et al., in prep



# Masses


- High BH masses because of **direct collapse** at low metallicity



Ziosi et al., in prep


# Masses

- High BH masses because of **direct collapse** at low metallicity
- **Chirp mass**  $m_{\text{chirp}} = \frac{(m_1 m_2)^{3/5}}{(m_1 + m_2)^{1/5}}$
- Why chirp mass:
  - $\nu_{\text{GW}} \propto m_{\text{chirp}}^{-5/8}$ ,  $h_{\text{GW}} \propto m_{\text{chirp}}^{5/3}$
  - So from observations we can reconstruct  $m_{\text{chirp}}$
  - In our model  $m_{\text{chirp}}$  strongly depends on  $Z$   
 $\Rightarrow$  Z-dependent BH mass model test
- **But:** in black chirp mass distribution of the best merger-candidates



# Coalescence times


- Time to reach **SMA=0** considering only **GW emission**
- $t_{GW} \propto \frac{a^4(1-e^2)^{7/2}}{m_1 m_2 m_{tot}}$  (Peters, 1964)
- GW emission: **SMA shrink** and **orbit circularization**
- **Dynamical outlier**: signal **detectability** depends on  $e$  (Brown & Zimmerman, 2010)
- 7 DBHs with  $t_{GW} < 13$  Gyr (0 for  $Z=Z_{\odot}$ )
- 17 DNSs with  $t_{GW} < 13$  Gyr, 11 DNS mergers during the simulations



# Conclusions



**DCOBs** during mergers **emits GWs** likely to be detected in the near future

- **Metallicity** is important:
  - **Heavier BHs** form at low Z
  - They tend to form DBHBs at **early times**
  - and these binaries are **more stable**:  $\Rightarrow$  DBHBs lifetimes are longer at low Z
- **Dynamics** is important
  - Dynamics enhances the formation of DCOBs: **97% of DBHBs** come **from exchanges**
  - Dynamics **hardens** binaries and can modify the **eccentricity**  $\Rightarrow$  increase **detection probability**



# Further work

- Increase statistical sample
- Dependence on ICs parameters:
  - YSC mass from  $10^3 M_{\odot}$  to  $10^5 M_{\odot}$
  - concentration (i.e. central dimensionless potential  $W_0$ , from 3 to 9)
  - asymmetries (presence of initial sub-clumps)
  - primordial binary fraction (from 0 to 0.5)
- Multiple systems analysis (triples, ...)
- External tidal field



# Thank you!